



Persistent Classical Swine Fever infection in newborn piglets

Uttenthal, Åse; Lohse, Louise; Rasmussen, Thomas Bruun; Nielsen, Jens

Publication date:
2011

[Link back to DTU Orbit](#)

Citation (APA):

Uttenthal, Å., Lohse, L., Rasmussen, T. B., & Nielsen, J. (2011). *Persistent Classical Swine Fever infection in newborn piglets*. Abstract from 8th ESVV International Pestivirus Symposium, Hanover, Lower Saxony, Germany. <http://www.pestivirus2011.de/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Persistent Classical Swine Fever infection in newborn piglets

Åse Uttenthal¹, Louise Lohse¹, Thomas Bruun Rasmussen¹, Jens Nielsen¹

¹DTU-Vet, Lindholm, 4771-Kalvehave, Denmark

Pestiviruses are unique in their ability to cause persistent infection (PI) in pigs infected in utero. In cattle, PI calves play an important role in maintenance of bovine viral diarrhoea virus infection in the herd. In pigs, the occurrence of classical swine fever virus (CSFV) PI piglets is anticipated to be epidemiologically important.

To study the course of CSFV PI in pigs, four sows were infected with 2009-CSFV Lithuania between day 50-60 of gestation. The sows gave birth to 66 piglets of which 55 % were live-born. Out of these, thirty % were considered to be PI pigs, while 40% were acutely infected and seroconverted rapidly. The status of the remaining 30% is unclear. Both PI and acutely infected piglets occurred in the same litters.

All piglets were tested CSFV antibody-negative at birth if precolostral blood samples were available. PI piglets quickly lost the maternally derived antibodies as free antibodies were not detected in serum even though the sows had Virus neutralization titer (Vnt) titers of 100. Non-PI piglets were able to raise active immunity, since specific antibodies to CSFV stabilized at a mean Vnt titer of 200. While some PI piglets showed growth retardation as well as central nervous disturbances, several others developed normally without showing clinical symptoms. The correlations between clinical signs, virus isolation, antibody levels and detection of CSFV by quantitative RT-PCR will be compared for PI and acutely infected piglets.

Acknowledgement: The project was supported by Directorate for Food, Fisheries and Agri-Business in Denmark (grant 2007-776).